



# STD Recommended Queries

# Chlamydia (2011)

All races/ethnicities | Both sexes | 2011 | All age groups




## Rate per 100,000 among selected population



Legend classified according to 2011 data.

## National Data By Sex




**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.

**Data Source:** The Atlas presents chlamydia, gonorrhea, and primary & secondary syphilis case report data for 2000 to 2011 and early latent syphilis case report data for 2003-2011. Data consist of case count or rate per 100,000 by disease, year of diagnosis, reporting area (state or territory), age group, race/ethnicity, and sex. **For more information, see: [Interpreting STD Surveillance Data](#).**


**Notes:** The Atlas includes data from STD case reports from 54 areas (all 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands). **Cases** of a given STD refer to confirmed diagnoses during a given time period. **Rates** per 100,000 population were calculated for each STD. The population denominators used to compute rates for the 50 states and the District of Columbia were based on the National Center for Health Statistics (NCHS) bridged-race population counts for 2000-2010. Due to the lack of data from the U.S. Census Bureau and small cell sizes, rates are not calculated for dependent areas (except Puerto Rico for age and sex). **Cell suppression:** CDC follows the 1996 CSTE data re-release rules for STD data. For state-level analysis, data are suppressed when the numerator for a given state is 3 or less. When suppressed, data are only available as state totals and no demographic data are shown. **Chlamydia:** Trends in chlamydia morbidity reporting from many state and local jurisdictions are more reflective of changes in diagnostic, screening, and reporting practices than of actual trends in disease incidence. **Suggested citation:** [NCHHSTP Atlas](#)

# Chlamydia (2011)

All races/ethnicities | Female | 2011 | All age groups




## Rate per 100,000 among selected population



Legend classified according to 2011 data.

## National Data By Age Group




**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.

**Data Source:** The Atlas presents chlamydia, gonorrhea, and primary & secondary syphilis case report data for 2000 to 2011 and early latent syphilis case report data for 2003-2011. Data consist of case count or rate per 100,000 by disease, year of diagnosis, reporting area (state or territory), age group, race/ethnicity, and sex. **For more information, see: [Interpreting STD Surveillance Data](#).**

**Notes:** The Atlas includes data from STD case reports from 54 areas (all 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands). **Cases** of a given STD refer to confirmed diagnoses during a given time period. **Rates** per 100,000 population were calculated for each STD. The population denominators used to compute rates for the 50 states and the District of Columbia were based on the National Center for Health Statistics (NCHS) bridged-race population counts for 2000-2010. Due to the lack of data from the U.S. Census Bureau and small cell sizes, rates are not calculated for dependent areas (except Puerto Rico for age and sex). **Cell suppression:** CDC follows the 1996 CSTE data re-release rules for STD data. For state-level analysis, data are suppressed when the numerator for a given state is 3 or less. When suppressed, data are only available as state totals and no demographic data are shown. **Chlamydia:** Trends in chlamydia morbidity reporting from many state and local jurisdictions are more reflective of changes in diagnostic, screening, and reporting practices than of actual trends in disease incidence. **Suggested citation:** [NCHHSTP Atlas](#)

# Gonorrhea (2011)

All races/ethnicities | Both sexes | Change over time (2000-2011) | All age groups




## Rate per 100,000 among selected population



Legend classified according to 2000-2011 data.

## National Data By Year



**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.

**Data Source:** The Atlas presents chlamydia, gonorrhea, and primary & secondary syphilis case report data for 2000 to 2011 and early latent syphilis case report data for 2003-2011. Data consist of case count or rate per 100,000 by disease, year of diagnosis, reporting area (state or territory), age group, race/ethnicity, and sex. For more information, see: [Interpreting STD Surveillance Data](#).

**Notes:** The Atlas includes data from STD case reports from 54 areas (all 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands). **Cases** of a given STD refer to confirmed diagnoses during a given time period. **Rates** per 100,000 population were calculated for each STD. The population denominators used to compute rates for the 50 states and the District of Columbia were based on the National Center for Health Statistics (NCHS) bridged-race population counts for 2000-2010. Due to the lack of data from the U.S. Census Bureau and small cell sizes, rates are not calculated for dependent areas (except Puerto Rico for age and sex). **Cell suppression:** CDC follows the 1996 CSTE data re-release rules for STD data. For state-level analysis, data are suppressed when the numerator for a given state is 3 or less. When suppressed, data are only available as state totals and no demographic data are shown. **Chlamydia:** Trends in chlamydia morbidity reporting from many state and local jurisdictions are more reflective of changes in diagnostic, screening, and reporting practices than of actual trends in disease incidence. **Suggested citation:** [NCHHSTP Atlas](#)

Centers for Disease Control and Prevention


National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention






# Gonorrhea (2011)

Black/African American | Both sexes | 2011 | All age groups




## Rate per 100,000 among selected population



Legend classified according to 2011 data.

## National Data By Age Group




**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.

**Data Source:** The Atlas presents chlamydia, gonorrhea, and primary & secondary syphilis case report data for 2000 to 2011 and early latent syphilis case report data for 2003-2011. Data consist of case count or rate per 100,000 by disease, year of diagnosis, reporting area (state or territory), age group, race/ethnicity, and sex. **For more information, see: [Interpreting STD Surveillance Data](#).**

**Notes:** The Atlas includes data from STD case reports from 54 areas (all 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands). **Cases** of a given STD refer to confirmed diagnoses during a given time period. **Rates** per 100,000 population were calculated for each STD. The population denominators used to compute rates for the 50 states and the District of Columbia were based on the National Center for Health Statistics (NCHS) bridged-race population counts for 2000-2010. Due to the lack of data from the U.S. Census Bureau and small cell sizes, rates are not calculated for dependent areas (except Puerto Rico for age and sex). **Cell suppression:** CDC follows the 1996 CSTE data re-release rules for STD data. For state-level analysis, data are suppressed when the numerator for a given state is 3 or less. When suppressed, data are only available as state totals and no demographic data are shown. **Chlamydia:** Trends in chlamydia morbidity reporting from many state and local jurisdictions are more reflective of changes in diagnostic, screening, and reporting practices than of actual trends in disease incidence. **Suggested citation: [NCHHSTP Atlas](#)**

# Primary and Secondary Syphilis (2011)

All races/ethnicities | Male | 2011 | All age groups



Alaska



Hawaii



District of Columbia



Puerto Rico



American Samoa



Guam



N. Mariana Islands



U.S. Virgin Islands

## Rate per 100,000 among selected population

□ Data not available

2.6 - 5.1

10.1 - 55.6

□ Data suppressed


5.2 - 7.3

0.4 - 2.1

7.4 - 9.6

Legend classified according to 2011 data.

## National Data By Age Group



**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.

**Data Source:** The Atlas presents chlamydia, gonorrhea, and primary & secondary syphilis case report data for 2000 to 2011 and early latent syphilis case report data for 2003-2011. Data consist of case count or rate per 100,000 by disease, year of diagnosis, reporting area (state or territory), age group, race/ethnicity, and sex. **For more information, see:** [Interpreting STD Surveillance Data](#).

**Notes:** The Atlas includes data from STD case reports from 54 areas (all 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands). **Cases** of a given STD refer to confirmed diagnoses during a given time period. **Rates** per 100,000 population were calculated for each STD. The population denominators used to compute rates for the 50 states and the District of Columbia were based on the National Center for Health Statistics (NCHS) bridged-race population counts for 2000-2010. Due to the lack of data from the U.S. Census Bureau and small cell sizes, rates are not calculated for dependent areas (except Puerto Rico for age and sex). **Cell suppression:** CDC follows the 1996 CSTE data re-release rules for STD data. For state-level analysis, data are suppressed when the numerator for a given state is 3 or less. When suppressed, data are only available as state totals and no demographic data are shown. **Chlamydia:** Trends in chlamydia morbidity reporting from many state and local jurisdictions are more reflective of changes in diagnostic, screening, and reporting practices than of actual trends in disease incidence. **Suggested citation:** [NCHHSTP Atlas](#)


Centers for Disease Control and Prevention

National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention



# Primary and Secondary Syphilis (2011)

All races/ethnicities | Male | 2011 | 15-19




## Rate per 100,000 among selected population



Legend classified according to 2011 data.

## National Data By Race/ethnicity



**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.


**Data Source:** The Atlas presents chlamydia, gonorrhea, and primary & secondary syphilis case report data for 2000 to 2011 and early latent syphilis case report data for 2003-2011. Data consist of case count or rate per 100,000 by disease, year of diagnosis, reporting area (state or territory), age group, race/ethnicity, and sex. **For more information, see:** [Interpreting STD Surveillance Data](#).

**Notes:** The Atlas includes data from STD case reports from 54 areas (all 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands). **Cases** of a given STD refer to confirmed diagnoses during a given time period. **Rates** per 100,000 population were calculated for each STD. The population denominators used to compute rates for the 50 states and the District of Columbia were based on the National Center for Health Statistics (NCHS) bridged-race population counts for 2000-2010. Due to the lack of data from the U.S. Census Bureau and small cell sizes, rates are not calculated for dependent areas (except Puerto Rico for age and sex). **Cell suppression:** CDC follows the 1996 CSTE data re-release rules for STD data. For state-level analysis, data are suppressed when the numerator for a given state is 3 or less. When suppressed, data are only available as state totals and no demographic data are shown. **Chlamydia:** Trends in chlamydia morbidity reporting from many state and local jurisdictions are more reflective of changes in diagnostic, screening, and reporting practices than of actual trends in disease incidence. **Suggested citation:** [NCHHSTP Atlas](#)



# Primary and Secondary Syphilis (2011)

All races/ethnicities | Female | 2011 | All age groups




## Rate per 100,000 among selected population



Legend classified according to 2011 data.

## National Data By Race/ethnicity



**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.

**Data Source:** The Atlas presents chlamydia, gonorrhea, and primary & secondary syphilis case report data for 2000 to 2011 and early latent syphilis case report data for 2003-2011. Data consist of case count or rate per 100,000 by disease, year of diagnosis, reporting area (state or territory), age group, race/ethnicity, and sex. **For more information, see:** [Interpreting STD Surveillance Data](#).

**Notes:** The Atlas includes data from STD case reports from 54 areas (all 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands). **Cases** of a given STD refer to confirmed diagnoses during a given time period. **Rates** per 100,000 population were calculated for each STD. The population denominators used to compute rates for the 50 states and the District of Columbia were based on the National Center for Health Statistics (NCHS) bridged-race population counts for 2000-2010. Due to the lack of data from the U.S. Census Bureau and small cell sizes, rates are not calculated for dependent areas (except Puerto Rico for age and sex). **Cell suppression:** CDC follows the 1996 CSTE data re-release rules for STD data. For state-level analysis, data are suppressed when the numerator for a given state is 3 or less. When suppressed, data are only available as state totals and no demographic data are shown. **Chlamydia:** Trends in chlamydia morbidity reporting from many state and local jurisdictions are more reflective of changes in diagnostic, screening, and reporting practices than of actual trends in disease incidence. **Suggested citation:** [NCHHSTP Atlas](#)

Centers for Disease Control and Prevention


National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention






# Early Latent Syphilis (2011)

All races/ethnicities | Female | 2011 | All age groups




## Rate per 100,000 among selected population



Legend classified according to 2011 data.

## National Data By Age Group



**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.

**Data Source:** The Atlas presents chlamydia, gonorrhea, and primary & secondary syphilis case report data for 2000 to 2011 and early latent syphilis case report data for 2003-2011. Data consist of case count or rate per 100,000 by disease, year of diagnosis, reporting area (state or territory), age group, race/ethnicity, and sex. For more information, see: [Interpreting STD Surveillance Data](#).

**Notes:** The Atlas includes data from STD case reports from 54 areas (all 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands). Cases of a given STD refer to confirmed diagnoses during a given time period. Rates per 100,000 population were calculated for each STD. The population denominators used to compute rates for the 50 states and the District of Columbia were based on the National Center for Health Statistics (NCHS) bridged-race population counts for 2000-2010. Due to the lack of data from the U.S. Census Bureau and small cell sizes, rates are not calculated for dependent areas (except Puerto Rico for age and sex). **Cell suppression:** CDC follows the 1996 CSTE data re-release rules for STD data. For state-level analysis, data are suppressed when the numerator for a given state is 3 or less. When suppressed, data are only available as state totals and no demographic data are shown. **Chlamydia:** Trends in chlamydia morbidity reporting from many state and local jurisdictions are more reflective of changes in diagnostic, screening, and reporting practices than of actual trends in disease incidence. **Suggested citation:** NCHHSTP Atlas


Centers for Disease Control and Prevention

National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention



# Early Latent Syphilis (2011)

All races/ethnicities | Male | 2011 | All age groups




## Rate per 100,000 among selected population



Legend classified according to 2011 data.

## National Data By Age Group



**Disclaimer:** This is a user-generated report. The findings and conclusions are those of the user and do not necessarily represent the views of the CDC.

**Data Source:** The Atlas presents chlamydia, gonorrhea, and primary & secondary syphilis case report data for 2000 to 2011 and early latent syphilis case report data for 2003-2011. Data consist of case count or rate per 100,000 by disease, year of diagnosis, reporting area (state or territory), age group, race/ethnicity, and sex. **For more information, see:** [Interpreting STD Surveillance Data](#).

**Notes:** The Atlas includes data from STD case reports from 54 areas (all 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands). **Cases** of a given STD refer to confirmed diagnoses during a given time period. **Rates** per 100,000 population were calculated for each STD. The population denominators used to compute rates for the 50 states and the District of Columbia were based on the National Center for Health Statistics (NCHS) bridged-race population counts for 2000-2010. Due to the lack of data from the U.S. Census Bureau and small cell sizes, rates are not calculated for dependent areas (except Puerto Rico for age and sex). **Cell suppression:** CDC follows the 1996 CSTE data re-release rules for STD data. For state-level analysis, data are suppressed when the numerator for a given state is 3 or less. When suppressed, data are only available as state totals and no demographic data are shown. **Chlamydia:** Trends in chlamydia morbidity reporting from many state and local jurisdictions are more reflective of changes in diagnostic, screening, and reporting practices than of actual trends in disease incidence. **Suggested citation:** [NCHHSTP Atlas](#)